



## Digital Receipt

This receipt acknowledges that **Turnitin** received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Dewa Ayu Putu Sri Masyeni  
Assignment title: Article  
Submission title: Antiretroviral therapy substitution am...  
File name: CRC\_pdf\_antiretroviral.pdf  
File size: 391.91K  
Page count: 5  
Word count: 2,200  
Character count: 11,417  
Submission date: 04-Mar-2021 10:53PM (UTC+0700)  
Submission ID: 1524129089

*Medical Technology & Environmental Health: Abdullah, Widhiy 7 Abdullah (eds)  
© 2020 Taylor & Francis Group, London, ISBN, 978-0-367-86033-0*

### Antiretroviral therapy substitution among HIV/AIDS patients visiting Sanjiwani Hospital, Bali

S Masyeni<sup>1</sup>, I W A Sudarsana and I D G Wedha Asmara  
*Universitas Warmadewa, Denpasar, Indonesia*

**Abstract.** Substitution of long-term ART-exposed to HIV/AIDS patient is common due to the ART adverse event. The study aim is to describe the reason for ART substitution among HIV/AIDS patient at Sanjiwani Hospital Bali. A retrospective study was conducted to the medical records of HIV/AIDS patients at Sanjiwani hospital Bali, during 2006-2018. Clinical data retrieved from the medical records and presented as descriptive data. Over twelve years, 1112 HIV/AIDS patients evaluated in the study. The ART regimen were Zidovudine-based ART, Tenofovir-based ART, and Stavudine-based, for 12.2%, 87.3%, 0.5%, respectively. There were 2.2 % switching of ART during the study period. The most reason was anemia (48%), followed by reducing of kidney function (28%), allergic reaction (16%), and 4% of nausea and suspected failure to ART clinically. We highlight that anemia is the main reason for ART substitution among HIV/AIDS patients.

#### 1. Introduction

Antiretroviral therapy with Highly Active Antiretroviral Therapy (HAART) in HIV infection converted a fatal condition into a chronic and manageable illness. In resource-limited countries, the ART regimen mostly consist of a combination of two nucleoside analogue reverse transcriptase inhibitors (NRTIs) such as zidovudine (AZT), lamivudine (3TC), tenofovir disoproxil fumarate (TDF) and one of non-nucleoside reverse transcriptase inhibitors (NNRTI) such as nevirapine (NVP) or efavirenz (EFV) [1]. The mode of action of the NRTIs competes with the natural deoxynucleotides for incorporation into the growing viral DNA chain. Unlike the natural deoxynucleotides substrate, NRTIs lack a 3'-hydroxyl group on the deoxyribose moiety; hence, following incorporation of the NRTIs, the newly performed deoxynucleotide cannot form the next 5'-3' phosphodiester bond needed to extend the DNA chain [2]. The NRTIs triphosphate inhibits the function of polymerase- $\gamma$ , the enzyme responsible for mitochondrial DNA (mtDNA) replication; hence, the depletion of mtDNA is common among HIV-treated person [3-5].

Even though HAART can fruitfully defeat viral replication in the long term, it is not without substantial toxicity, which can extremely complicate treatment effectiveness. Central toxicity that has documented for more than a decade. The severity of the adverse events are range from mild to life-threatening with short and long term effects is NRTI-related mitochondrial toxicity, which exhibits as severe side effects such as hepatic failure, cardiac dysfunction, skeletal myopathies and lactic acidosis [6]. Adverse event of ART reported as high as 54% on AZT in which the most ordinary adverse events were a pain (30%) and skin rashes (18%) [7]. The general principle of ART toxicities is depended on the severity of the adverse events. Mild toxicities do not require termination of therapy or drug

# Antiretroviral therapy substitution among HIV/AIDS patients visiting Sanjiwani Hospital, Bali

*by* Dewa Ayu Putu Sri Masyeni

---

**Submission date:** 04-Mar-2021 07:53AM (UTC-0800)

**Submission ID:** 1524129089

**File name:** CRC\_pdf\_antiretroviral.pdf (391.91K)

**Word count:** 2200

**Character count:** 11417

## Antiretroviral therapy substitution among HIV/AIDS patients visiting Sanjiwani Hospital, Bali

S Masyeni\*, I W A Sudiarsana and I D G Wedha Asmara

Universitas Warmadewa, Denpasar, Indonesia

**Abstract.** Substitution of long-term ART-exposed to HIV/AIDS patient is common due to the ART adverse event. The study aim is to describe the reason for ART substitution among HIV/AIDS patient at Sanjiwani Hospital Bali. A retrospective study was conducted to the medical records of HIV/AIDS patients at Sanjiwani hospital Bali, during 2006-2018. Clinical data retrieved from the medical records and presented as descriptive data. Over twelve years, 1112 HIV/AIDS patients evaluated in the study. The ART regimen were Zidovudine-based ART, Tenofovir-based ART, and Stavudine-based, for 12.2%, 87.3%, 0.5%, respectively. There were 2.2 % switching of ART during the study period. The most reason was anemia (48%), followed by reducing of kidney function (28%), allergic reaction (16%), and 4% of nausea and suspected failure to ART clinically. We highlight that anemia is the main reason for ART substitution among HIV/AIDS patients.

### 1. Introduction

Antiretroviral therapy with Highly Active Antiretroviral Therapy (HAART) in HIV infection converted a fatal condition into a chronic and manageable illness. In resource-limited countries, the ART regimen mostly consist of a combination of two nucleoside analogue reverse transcriptase inhibitors (NRTIs) such as zidovudine (AZT), lamivudine (3TC), tenofovir disoproxil fumarate (TDF) and one of non-nucleoside reverse transcriptase inhibitors (NNRTI) such as nevirapine (NVP) or efavirenz (EFV) [1]. The mode of action of the NRTIs competes with the natural deoxynucleotides for incorporation into the growing viral DNA chain. Unlike the natural deoxynucleotides substrate, NRTIs lack a 3'-hydroxyl group on the deoxyribose moiety; hence, following incorporation of the NRTIs, the newly performed deoxynucleotide cannot form the next 5'-3' phosphodiester bond needed to extend the DNA chain [2]. The NRTIs triphosphate inhibits the function of polymerase- $\gamma$ , the enzyme responsible for mitochondrial DNA (mtDNA) replication; hence, the depletion of mtDNA is common among HIV-treated person [3-5].

Even though HAART can fruitfully defeat viral replication in the long term, it is not without substantial toxicity, which can extremely conciliate treatment effectiveness. Central toxicity that has documented for more than a decade. The severity of the adverse events are range from mild to life-threatening with short and long term effects is NRTI-related mitochondrial toxicity, which exhibits as severe side effects such as hepatic failure, cardiac dysfunction, skeletal myopathies and lactic acidosis [6]. Adverse event of ART reported as high as 54% on AZT, in which the most ordinary adverse events were a pain (30%) and skin rashes (18%) [7]. The general principle of ART toxicities is depended on the severity of the adverse events. Mild toxicities do not require termination of therapy or drug

substitution, and symptomatic management may give (e.g., antihistamines for a mild rash). Moderate or severe toxicities may require substitution with a drug in the same ART class but with a different toxicity profile, or with a drug in a different class, but do not require discontinuation of all ART. Severe life-threatening toxicities need cessation of all ARV drugs, and the commencement of proper supportive therapy until the patient alleviated, and the toxicity is committed. Substitution of long-term ART-exposed to HIV/AIDS patients are common due to the ART adverse event [7].

However, only a little information is known about ART adverse events in many HIV programs in the public health sector of developing countries. The study aim is to describe the reason for ART substitution among HIV/AIDS patient at Sanjiwani Hospital Bali.

## 2. Methods

The current study was a hospital-based retrospective observational study conducted at HIV care clinics in Gianyar Bali from 2006- 2018. The hospital has an HIV clinic, staffed with health professionals trained in ART treatment and adherence counseling services. Clinical data retrieved from the medical records and presented as descriptive data. The reasons for substitution retrieved from the medical record. A data-gathering format used to collect data on the demographic settings, the starting and changing regimens, the period of the initial therapy, CD4 count, and World Health Organization (WHO) stage of the disease, and reasons for regimen substitution. Adverse drug reactions (ADR) is defined as the occurrence of adverse events such as diarrhea, nausea, vomiting, anemia, rash, fatigue, peripheral neuropathy, lipodystrophy, metabolic disturbances or any other related to HAART. Substitution defined as single or triple drug change due to side effects and initiating another drug of the same class and or another category.

## 3. Results


A total of 1.094 medical records of HIV-infected at Sanjiwani Hospital assessed in the study. Female patients account in the minority, as high as 29% and 7(0.6%) of the female participants was pregnant. The median age of the participants is 32.5 (IQR 13) years old. Total CD4+  $\leq 100$  cell/mm<sup>3</sup>, CD4+ 101-200 cell/mm<sup>3</sup>, CD4+ 201-350 cell/mm<sup>3</sup>, CD4+ 351-499 cell/mm<sup>3</sup>, and CD4+  $\geq 500$  cell/ml accounting for 347 (31.7%), 139 (12.7%), 173 (15.8%) and 37 (93.4%) respectively.

**Table 1.** Characteristic of the participants (N=1094)

Variable	Frequency	Percentage (%)
Gender		
Male	777	71
Age		
18-30	470	42.96
31-40	382	34.92
41-50	166	15.17
51-60	58	5.31
61-70	17	1.55
>70	1	0.09
Initial Body Weight		
Median (IQR)	55 (12)	
Initial CD4 (IQR)	136 (246)	
HIV stage (WHO)		
Stage 1	115	10.5
Stage 2	109	10
Stage 3	431	39.4
Stage 4	439	40.1

The first line original ART consists of stavudine-based, tenofovir-based, and zidovudine-based ART. The stavudine ART in combination with lamivudine and nevirapine or efavirenz, as well as the combination of TDF and ZDV. Characteristic of the participants presented in table 1. A majority of the patients were on the combination of tenofovir+lamivudine+efavirenz, the fixed-dose combination ART (738/67.5%).

Total switching of the ART found in 26 (2.37%) patients. Ten of 26 (38.46) switching was on ZDV-based as the original ART. Adverse events of ART found in 22 (2.01%) patients. Type of adverse events is anemia 4 (18.18%), itchy 10 (45.45%), a decrease of kidney function seven (31.82) and gastrointestinal problem 1 (4.55%). The adverse effect anemia and itchy mostly relate to the combination of ZDV+3TC+NVP and another first-line ART, TDF+3TC+EFV is the most ART of choice substitute the previous ART. On the other hand, reducing kidney function may relate to TDF-based ART and change to ZDV-based ART. Fifty percent of the switching occurs in the first year of treatment.



**Figure 1.** Profile of ART among the HIV-infected patients

The switching to second-line ART found in four (0.36%) of the patients due to the suspicion of treatment failure whether clinical failure 33.3% and immunological failure 66.7%. The ART of choice for switching due to treatment failure is boosted lopinavir/ritonavir in combination with TDF+3TC.

#### 4. Discussion

Acquired immunodeficiency syndrome (AIDS) caused by the human immunodeficiency virus (HIV) is a major global health problem [8]. There are several studies of HIV infection in Bali due to subtype characteristic [9], toxicity [5], adherence [10], and co-infection [11] which reflect the epidemic of HIV infection in Bali. The main reasons for treatment conversion might be due to, adverse events, poor adherence, a desire for pregnancy, treatment failure [12]. The high finding of male patients who need ART substitution in the current study is supported by another study where they found as high as 53.7%.

The median CD4 of the study participants is 136 cell/mm<sup>3</sup>. In contrast with the finding of the previous study where they found the median CD4 of the patients was 201 cell/mm<sup>3</sup> [13]. The discrepancy may associate with the social demographics of the countries affect the immune status of the patients. The study found toxicity (88%) as the most reason to change the ART regimen. A concordance finding reported by other study that found as high as 72.73% ART changing due to toxicity [14]. This similar finding may due to the HIV sub-type is HIV-1, but we do not assess the genetic diversity of the patients.

The most common primary ART in the current study is TDF-based regimen, which is in contrast with previous study Zidovudine-based ART where the primary ART was ZDV-based ART [15]. In Indonesia, the availability of TDF since 2014 become the ART of choice due to merely way to replace others ART of choice. Likewise, the ZDV adverse events such as anemia may associate with the common ART to switch to another regimen in this study. Another study found ART substitution most commonly found due to NVP toxicities [16]. This study found toxicity due to NVP although in the minority, that may explain with the most ART use at the hospital is TDF+3TC+EFV.

## 5. Conclusion

We highlight the most ART substitution in the study due to ART toxicities, instead of treatment failure. This finding may help the physician in monitoring the ART adverse event in improving the services for the patient's convenience.

## 6. Acknowledgment

We thank the honorable doctor, nurse, and Faculty of Medicine and Health Sciences Universitas Warmadewa for the great support of this work

## 7. Conflict of interest

The authors declare no conflict of interest

## 8. References

- [1] Kementerian Kesehatan Republik Indonesia 2014 Pedoman pengobatan antiretroviral *Peratur. Menteri Kesehat. Republik Indones. Nomor 87 Tahun 2014* pp 1–121
- [2] Tressler R and Godfrey C 2012 NRTI backbone in HIV treatment *Drugs* **72(16)** pp 2051–62
- [3] Wagner T A, Lin C-H, Tobin N H, Côté H C F, Sloan D D and Jerome K R 2013 Quantification of mitochondrial toxicity in HIV-infected individuals by quantitative PCR compared to flow cytometry *Cytom. Part B Clin. Cytom. [Internet]* **84B (1)** pp 55–8 Available from: <http://doi.wiley.com/10.1002/cyto.b.21045>
- [4] Montaner J S G, Côté H C F, Harris M, Hogg R S, Yip B and Harrigan P R 2004 Nucleoside-related mitochondrial toxicity among HIV-infected patients receiving antiretroviral therapy: insights from the evaluation of venous lactic acid and peripheral blood mitochondrial DNA *Clin. Infect. Dis. [Internet]* **38 Suppl. 2** pp S73–9 Available from: <http://www.ncbi.nlm.nih.gov/pubmed/14986278>
- [5] Masyeni S, Sintya E, Megawati D, Sukmawati N M H, Budiya D G and Aryastuti S A 2018 HIV Evaluation of antiretroviral effect on mitochondrial DNA depletion among HIV-infected patients in Bali *AIDS (Auckl)* **10** pp 145–50
- [6] Gudina E K, Teklu A M, Berhan A, Gebreegziabhier A, Seyoum T and Nega A 2017 Magnitude of antiretroviral drug toxicity in adult HIV patients in Ethiopia: A cohort study at seven teaching hospitals *Ethiop. J. Health Sci. [Internet]* **Suppl. 1** pp 39–52 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28465652>
- [7] Eluwa G I, Badru T, Agu K A, Akpoigbe K J, Chabikuli O and Hamelmann C 2012 Adverse drug reactions to antiretroviral therapy (ARVs): incidence, type and risk factors in Nigeria *BMC Clin. Pharmacol.* **12** p 7
- [8] Van Sighem A, Nakagawa F, De Angelis D, Quinten C, Bezemer D and De Coul E O 2015 Estimating HIV incidence, time to diagnosis, and the undiagnosed HIV epidemic using routine surveillance data *Epidemiology* **26(5)** pp 653–60
- [9] Khairunisa S Q, Masyeni S, Witaningrum A M, Indriati D W, Kotaki T and Ueda S 2018 Genotypic characterization of human immunodeficiency virus type 1 isolated in Bali, Indonesia in 2016 *HIV AIDS Rev. Int. J. HIV-Related Probl.* **17(2)** pp 81–90.
- [10] Jiamsakul A, Kumarasamy N, Ditangco R, Li P C K, Phanuphak P, Sirisanthana T, et al 2014 *J. Int. AIDS Soc. Factors associated with suboptimal adherence to antiretroviral therapy in Asia*

**17(1)** p 18911

- [11] Juliari I G A M and Susila N K N 2018. Occular syphilis in HIV-positive male. *IOP Conference Series: Material Sciences and Engineering*. **434(1)** 012340
- [12] Haile G S and Berha A B 2019 Predictors of treatment failure, time to switch and reasons for switching to second line antiretroviral therapy in HIV infected children receiving first line antiretroviral therapy at a tertiary care hospital in Ethiopia *BMC Pediatr*. **19(1)** p 37
- [12] Zhang F, Dou Z, Ma Y, Zhang Y, Zhao Y and Zhao D 2011 Effect of earlier initiation of antiretroviral treatment and increased treatment coverage on HIV-related mortality in China: a national observational cohort study *Lancet Infect. Dis.* **11(7)** pp 516–24
- [13] Assefa D and Hussein N 2014 Reasons for regimen change among HIV/AIDS patients initiated on first line highly active antiretroviral therapy in Fitche Hospital, Oromia, Ethiopia *Adv. Pharmacol. Pharm.* **2(5)** pp 77–83
- [14] Sandeep B, Chavan VR, Raghunandan M, Arshad M and Sayana S B 2014 Factors influencing the substitution of antiretroviral therapy in human immunodeficiency virus/acquired immunodeficiency syndrome patients on first line highly active antiretroviral therapy *Asian J. Pharm. Clin. Res.* **7 (5)** pp 117–20
- [15] Boulle A, Orrell C, Kaplan R, Van Cutsem G, McNally M and Hilderbrand K 2007 Substitutions due to antiretroviral toxicity or contraindication in the first 3 years of antiretroviral therapy in a large South African cohort



# Antiretroviral therapy substitution among HIV/AIDS patients visiting Sanjiwani Hospital, Bali

## ORIGINALITY REPORT

20%

SIMILARITY INDEX

16%

INTERNET SOURCES

16%

PUBLICATIONS

10%

STUDENT PAPERS

## PRIMARY SOURCES

1

[whqlibdoc.who.int](http://whqlibdoc.who.int)

Internet Source

5%

2

Submitted to University of the Highlands and Islands Millennium Institute

Student Paper

3%

3

François-René Alexandre, Rachid Rahali, Houcine Rahali, Sandra Guillon et al. "Synthesis and Antiviral Evaluation of Carbocyclic Nucleoside Analogs of Nucleoside Reverse Transcriptase Translocation Inhibitor MK-8591 (4'-Ethynyl-2-fluoro-2'-deoxyadenosine)", Journal of Medicinal Chemistry, 2018

Publication

2%

4

[link.springer.com](http://link.springer.com)

Internet Source

2%

5

"Encyclopedia of AIDS", Springer Science and Business Media LLC, 2018

Publication

1%

[patua.iec.gov.br](http://patua.iec.gov.br)



6

Internet Source

1%

7

"Assessment of adverse drug reactions to antiretroviral agents among HIV patients",  
Journal of Applied Pharmaceutical Science,  
2018

Publication

1%

8

[www.intmedpress.com](http://www.intmedpress.com)

Internet Source

1%

9

[www.civilejournal.org](http://www.civilejournal.org)

Internet Source

1%

10

Getiye Dejenu Kibret, Tadesse Awoke Ayele,  
Adino Tesfahun. "Incidence and Predictors of  
Sever Adverse Drug Reactions Among Patients  
on Antiretroviral Therapy at Debre Markos  
Referral Hospital, Northwest Ethiopia",  
Research Square, 2019

Publication

1%

11

[www.upo.es](http://www.upo.es)

Internet Source

1%

12

[bmccresnotes.biomedcentral.com](http://bmccresnotes.biomedcentral.com)

Internet Source

&lt;1%

13

[springerplus.springeropen.com](http://springerplus.springeropen.com)

Internet Source

&lt;1%

14

[humanhealth-sumit.blogspot.com](http://humanhealth-sumit.blogspot.com)

<1 %

15

Sri Masyeni, Erly Sintya, Dewi Megawati, Ni Made Hegard Sukmawati et al. "Evaluation of antiretroviral effect on mitochondrial DNA depletion among HIV-infected patients in Bali", HIV/AIDS - Research and Palliative Care, 2018

Publication

<1 %

16

[www.sinta.ristekbrin.go.id](http://www.sinta.ristekbrin.go.id)

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On